

LETTER TO THE EDITOR

Spurious oscillations from local self-interaction correction in high energy photoionization calculations for metal clusters

M.E. Madjet, Himadri S. Chakraborty § , and Jan-M. Rost

Max-Planck-Institut für Physik Komplexer Systeme, Nöthnitzer Strasse 38, D-01187 Dresden, Germany

Abstract. We find that for simple metal clusters a single-electron description of the ground state employing self-interaction correction (SIC) in the framework of local-density approximation strongly contaminates the high energy photoionization cross sections with spurious oscillations for a subshell containing node(s). This effect is shown connected to the unphysical structure that SIC generates in ensuing state-dependent radial potentials around a position where the respective orbital density attains nodal zero. Non-local Hartree-Fock that *exactly* eliminates the electron self-interaction is found entirely free from this effect. It is inferred that while SIC is largely unimportant in high photon-energies, any implementation of it within the local frame can induce unphysical oscillations in the high energy photospectra of metal clusters pointing to a general need for caution in choosing appropriate theoretical tools.

PACS numbers: 31.15.Ew, 36.40.Cg, 36.40.Vz

The local-density approximation (LDA), along with its time-dependent version, is a standard theoretical technique to describe the structure and dynamics of large systems. From a practical standpoint, LDA is typically preferred to other conventional many-body methods (such as, Hartree-Fock (HF) or techniques based on configuration-interactions) because of its relatively low computational costs. In the context of the studies involving static and dynamical properties of simple metal clusters LDA has proved to be particularly successful [1-3]. However, a well known drawback of LDA is that it only partially accounts for unphysical electron self-interactions. As a consequence, the resulting potential for a finite system decays exponentially at large distance instead of producing the physical $1/r$ behavior. To render the long distance behavior of the LDA potential realistic, therefore, approximation schemes have been suggested [4]. The most general and widely applied to remedy the error is the one proposed by Perdew and Zunger [5], which concerns an orbit-by-orbit elimination of self-interaction, although the scheme immediately makes the potential state-dependent. The self-interaction corrected LDA (LDA-SIC) improves remarkably the vast variety of results related to many structural properties of physical systems: for instance,

§ To whom correspondence should be addressed (himadri@mpipks-dresden.mpg.de)

improvements in total energies of atoms, allowance for self-consistent bound solutions for negative ions, prediction of orbital energies that are close to electron removal energies thus restoring Koopmans' theorem, ensuring dissociation of heteronuclear molecules to neutral fragments, improvement of the band gap in solids etc. (a good account in this regard may be found through Ref. 4). In the dynamical regime too, especially in the context of low-energy photoionization of simple metal clusters, the description of the electronic ground state via LDA-SIC results in important many-body effects including single electron Rydberg resonances [6,7].

At photon-energies well beyond the ionization threshold the photospectrum shows special qualitative behavior. For spherical jellium clusters over this energy range theory predicts a characteristic oscillation in the cross section with a frequency connected to the cluster diameter. The mechanism behind this oscillatory pattern is the interference of electron waves emanated from equivalent sites of the cluster edge [8]. While there has been no experimental study on metal clusters, oscillations in the photoelectron intensity are indeed observed for fullerene molecules [9]. Generically, the high energy photoionization process should be rather sensitive to the degree of accuracy in the description of the ground state. This can be understood from the fact that in an independent particle model the high-energy transition matrix element has a leading contribution from the Fourier transform of the ground state wavefunction to the photoelectron momentum space (or retarded-momentum space if non-dipole interactions are included) [10]. From such an elementary viewpoint, LDA-SIC may also appear to be a suitable tool for the high energy photoionization studies of various cluster systems. However, this paper shows that while the correction for self-interaction is virtually unimportant in the study of energetic photoionization of metal clusters, any approximation to it in a local frame can generate spurious oscillations in the cross section for photoelectrons emerging from subshells having orbital node(s). The point is illustrated by presenting calculations on Na_{20} , which can be well described by a spherical jellium model, and which is the smallest system ($1s^2 1p^6 1d^{10} 2s^2$) having one subshell ($2s$) with a node.

The usual single electron potential in the Kohn-Sham LDA formalism is

$$V_{\text{KS}}(\vec{r}) = V_{\text{jel}}(\vec{r}) + V_{\text{d}}[\rho(\vec{r})] + V_{\text{xc}}[\rho(\vec{r})] \quad (1)$$

where the terms on the right-hand-side are respectively jellium, direct (Hartree) $V_{\text{d}}[\rho(\vec{r})] = \int d\vec{r}' \rho(\vec{r}')/|\vec{r} - \vec{r}'|$, and exchange-correlation potentials. The ground state electronic density $\rho(\vec{r})$ is defined in terms of single-electron densities ρ_i and orbitals ϕ_i :

$$\rho(\vec{r}) = \sum_{i=1}^N \rho_i(\vec{r}) = \sum_i |\phi_i(\vec{r})|^2$$

As mentioned earlier, an approximate prescription for SIC to this LDA potential (1) is to eliminate orbitalwise from the outset those terms which represent an electron i interacting to itself. The resulting orbital-specific potentials, therefore, are

$$V_{\text{SIC}}^i(\vec{r}) = V_{\text{jel}}(\vec{r}) + \int d\vec{r}' \frac{\rho(\vec{r}') - \rho_i(\vec{r}')}{|\vec{r} - \vec{r}'|} + V_{\text{xc}}[\rho(\vec{r})] - V_{\text{xc}}[\rho_i(\vec{r})] \quad (2)$$

As the exact form of V_{xc} is unknown a widely used scheme is to employ the formula[11]:

$$V_{xc}[\rho(\vec{r})] = - \left(\frac{3\rho(\vec{r})}{\pi} \right)^{1/3} - 0.0333 \log \left[1 + 11.4 \left(\frac{4\pi\rho(\vec{r})}{3} \right)^{1/3} \right] \quad (3)$$

The first term on the right-hand-side in the above expression is exactly derivable by a variational approach from the HF exchange energy of a uniform electron system with a uniform positively charged background; the second term is the so called correlation potential, a quantity not borne in HF formalism. We use LDA potentials both with and without SIC approximation to calculate the dipole photoionization cross sections upto approximately 1 KeV photon-energy for each subshell of the Na_{20} cluster in the independent particle frame[†]. Calculations are also performed in the self-consistent HF scheme to better identify the origin of the resulting discrepancy between the two LDA predictions. Quantities are in atomic units throughout, except where specified otherwise.

LDA and LDA-SIC cross sections for each of the $1s$, $1p$ and $1d$ subshells are found to be almost identical at high enough energies showing a single monotonic oscillation. Results using HF for these subshells yield similar qualitative behavior. The situation, however, is quite different for the $2s$ photoionization. Figure 1 presents $2s$ cross sections as obtained through LDA, LDA-SIC, and HF, as a function of $2s$ photoelectron momentum $k_{2s} = \sqrt{2(E - I_{2s})}$, with I_{2s} (~ 3.5 eV) being the $2s$ ionization threshold. Generally, in the low-energy range for all subshells of Na_{20} HF predictions are different from LDA owing to the partly non-identical ground state correlation they account for and this causes a constant phase difference between them at higher energies, where such correlation effects are insignificant. Bearing this in mind we find in figure 1 that while LDA and HF again maintain the same trend oscillationwise, LDA-SIC points to a progressively strong qualitative difference starting roughly from 40 eV photon-energy. To identify closely the discrepancy between σ_{2s} with and without SIC we have evaluated the Fourier transforms of $\sigma_{2s}(k_{2s})$ (see figure 2). Both LDA and HF are seen to have approximately the same Fourier spectrum with just one peak. But LDA-SIC contains three additional peaks beside the one that is common to all three spectra. This common frequency is connected to the diameter of the cluster. In fact, a simple theoretical analysis shows that high energy photo cross sections of a spherical jellium cluster oscillate in the respective photoelectron momentum space at a frequency $2R_C$, where R_C is the cluster radius [8]. However, where do the other frequencies in the LDA-SIC $2s$ cross section come from?

In order to answer this we need to take a close look at the single-electron ground state LDA and LDA-SIC radial potentials. As pointed out earlier, in LDA formalism “all” electrons of the system in the ground state experience the same potential defined by equation (1) which for Na_{20} is denoted by the dotted curve in figure 3. The potential, as is typical for a cluster, is flat in the interior region (region of de-localized quasi-free electrons) while showing a strong screening at the edge around R_C ; the unphysical

[†] Of course at such high energy the Na^+ core will ionize. However, the inclusion of this effect, going beyond the jellium frame, will not change our result qualitatively.

exponential decay at the long range may be noted. Switching to the LDA-SIC scheme, electrons in every orbital now feel a distinctly different potential (see equation (2)) with an approximately correct long range behavior as represented by four solid curves in figure 3. In this group of four SIC potentials the ones for $1s$, $1p$, and $1d$ look qualitatively similar to the LDA potential but are slightly deeper. The $2s$ potential, on the other hand, exhibits a unique feature: a strong local variation around the position $r = R_n$. To pin down how this structure in the $2s$ LDA-SIC potential comes about we need to focus on the SIC exchange correction $V_{xc}[\rho_{2s}(\vec{r})]$. This quantity, with reference to expression (3), can be explicitly written as:

$$V_{xc}[\rho_{2s}(\vec{r})] = - \left(\frac{3\rho_{2s}(\vec{r})}{\pi} \right)^{1/3} - 0.0333 \log \left[1 + 11.4 \left(\frac{4\pi\rho_{2s}(\vec{r})}{3} \right)^{1/3} \right] \quad (4)$$

The $2s$ orbital density, $\rho_{2s}(\vec{r}) = |\phi_{2s}(\vec{r})|^2$, in the above equation, vanishes at $r = R_n$ as the $2s$ radial wavefunction passes through its node at R_n . Consequently, $V_{xc}[\rho_{2s}(\vec{r})]$ generates a cusp-like structure in the neighborhood of R_n that shows up in the LDA-SIC potential profile for the $2s$ orbital. Since the behavior here is directly connected to the zero in the $2s$ electron density we stress that it must also occur in any alternate prescription for $V_{xc}[\rho_{2s}(\vec{r})]$ different from formula (2). We further emphasize that this structure is entirely an artifact of an externally imposed SIC in a purely local frame, which certainly is an approximation since a complete cancellation of self-interactions requires an appropriate non-local treatment of the electron-exchange phenomenon as in the HF formalism. In fact, a forced localization of the exchange (Fock) term in the HF scheme does indeed produce an infinite singularity in the potential at the zero of the corresponding one-electron state function [12]. Nevertheless, the structure from LDA-SIC has a direct bearing on the subsequent $2s$ photoionization matrix element by producing an unphysical oscillation.

To behold the underlying mechanism let us consider the photoionization dipole matrix element. We use for convenience the acceleration gauge representation of the dipole interaction that involves the gradient of the potential seen by the outgoing electron. After carrying out angular integration with the assumptions of spherical symmetry and unpolarized light, one is left with a reduced radial matrix element for a dipole transition $nl \rightarrow \epsilon l'$ that in the acceleration formalism is $\langle \psi_{\epsilon l'} | dV/dr | \psi_{nl} \rangle$. Figure 4 shows that the derivatives of both the LDA potential and the LDA-SIC $2s$ potential peak close to $r = R_c$. In fact, the first derivative of any general cluster potential always peaks at the edge R_c , and therefore, the overlap integral in the radial matrix element has dominating contribution coming from the edge [8]. Further, for high enough energy $\psi_{\epsilon l'}$ can be described in the first Born picture as a spherical wave with asymptotic form $\cos(k_{nl}r + \delta_{l'})$. This immediately suggests that the matrix element will oscillate in the k_{nl} space with roughly a frequency that is equal to the distance of the peak derivative point from the origin. As a consequence, resulting cross sections should exhibit an oscillation as a function of k_{nl} with a frequency $2R_c$ (since the cross section is the squared modulus of the matrix element). As mentioned before, this

effect is already known and can be related to the common frequency peak in figure 2. However, something additional happens for the LDA-SIC case. The structure induced by the wavefunction node in the LDA-SIC potential for the $2s$ orbital produces a sharp discontinuity at R_n in dV_{2s}/dr , as also seen in figure 4. Such a derivative-discontinuity induces a second oscillation in the respective overlap integral with a frequency about R_n [13]. Subsequently, the $2s$ cross section with SIC acquires four oscillation frequencies: $R_c - R_n$, $2R_n$, $R_c + R_n$, and $2R_c$ (see figure 2) as a result of the interference. Evidently, the first three frequencies are artificial being connected to the unphysical structure in the potential. Non-local HF, which exactly eliminates electron self-interaction terms, is free from this effect (as is seen from figures 1 and 2). Moreover, the qualitative agreement of HF with LDA suggests that SIC is practically unimportant at high enough photon-energy. This is simply because with predominant contribution coming from the potential edge for higher energies any improvement in the asymptotic behavior of the wavefunction does not significantly influence the overlap integral. Therefore, for large systems where HF becomes computationally impracticable the usual LDA may be a safe choice in the high-energy regime. On the other hand, the fact that slower photoelectrons with their longer wavelength can hardly “resolve” this nodal structure explains why low-energy cross sections in LDA-SIC are practically uncontaminated. It is also simple to understand that there is nothing special about $2s$ photoelectrons, for the effect must also be present in the case of subshells having more than one node.

The characteristic potential for any de-localized electron system, as in a metal cluster, has a nearly flat interior region. Any rapid variation in this potential occurring in a small range can, therefore, have considerable effect on the photoionization overlap integral by significantly altering the amplitude of the continuum wave across this range. For atomic systems, however, electrons are far more localized owing to the strong nuclear attraction, and therefore, wavefunctions are far more compact around the nucleus. The near-Coulombic shape of a typical atomic potential with steep slope close to the origin can practically overwhelm any local variation as the one discussed in this paper. In order to verify this, we applied LDA-SIC for some typical cases of atomic photoionization without any problem.

It is true that SIC in the LDA frame induces certain extensiveness in the calculations by making the potential state-dependent. One possible simplification is to average over all such state-dependent potentials and use the averaged one for all electrons. We applied such an average-SIC potential to examine whether or not the effect reduces. We find that not only the effect survives but that it also now substantially affects photoelectrons from subshells without a node because the wavefunction overlap across the nodal zone is rather strong for them since their ground state wavefunctions are large in this region. However, it remains to be seen what happens if the potential is further approximated by a simplified-implementation of SIC, namely, the optimized effective potential method [14]. Finally, it has recently been found in the context of atoms that the independent particle model breaks down for the high energy photoionization due to the interchannel coupling effect [15]. There is no *a priori* reason to assume that this will not be the case

for cluster systems, although no study has yet been made. Nevertheless, in the future even if a multi-channel frame (namely, the time-dependent LDA which is akin to the random-phase approximation) is needed to characterize the energetic photoionization of clusters, this spurious effect will remain, at least qualitatively, and may also affect those channels whose single channel description is otherwise error-free.

To summarize, we have shown that the theoretical analysis in the framework of LDA with SIC incorporated may invoke unphysical strong qualitative variations in high energy photospectra of metal clusters for electrons emitted from a subshell with node(s); although there is no denying that LDA-SIC is one of the strong methodologies available to address low-energy processes. Through a comparison with the results via non-local HF, that is intrinsically free from the self-interaction error, we conclude that the difficulty is connected to an inexact footing of SIC in the LDA formalism. Hence, it is important to choose appropriate theoretical techniques suitable for a given energy range to avoid mis-interpretation of various effects in cluster photo-dynamical studies.

We thank Professor Steven T. Manson of GSU-Atlanta, USA, for making useful comments on the manuscript.

References

- [1] Calvayrac F, Reinhard P -G, Suraud E, and Ullrich C A 2000 *Phys. Rep.* **337** 493
- [2] 1999 *Metal Clusters*, edited by Ekardt W (New York: Wiley)
- [3] Brack M 1993 *Rev. Mod. Phys.* **65** 677
- [4] Perdew J P and Ernzerhof 1998 *Electronic Density Functional Theory Recent Progress and New Directions*, edited by Dobson John F, Vignale Giovanni and Das Mukunda P (New York: Plenum Press) p 31
- [5] Perdew J P and Zunger A 1981 *Phys. Rev. B* **23** 5048
- [6] Madjet M E and Hervieux P A 1999 *European Phys. J. D* **9** 217
- [7] Pacheco J M and Ekardt W 1992 *Z. Phys. D* **24** 65
- [8] Frank Olaf and Rost Jan M 1996 *Z. Phys. D* **38** 59; 1997 *Chem. Phys. Letts.* **271** 367
- [9] Xu Y B, Tan M Q, and Becker U 1996 *Phys. Rev. Letts.* **76** 3538; Liebsch T, Hentges R, Rüdél A, Viehhaus J, Becker U, and Schlögl R 1997 *Chem. Phys. Letts.* **279** 197; Becker Uwe, Gessner Oliver and Rüdél Andy 2000 *J. Elec. Spect. Rel. Phen* **108** 189
- [10] Bethe Hans A and Salpeter Edwin E, in *Quantum Mechanics of One- and Two-Electron Atoms* (Plenum) 1977, p 299.
- [11] Gunnerson O and Lundqvist B I 1976 *Phys. Rev. B* **13** 4274
- [12] Hansen M and Nishioka H 1993 *Z. Phys. D* **28** 73
- [13] Oscillation in the cross section from derivative discontinuity in the single electron potential is known in the context of atomic photoionization. Ref: Amusia M Ya, Band I M, Ivandov V K, Kupchenko V A, and Trzhashovskaya M B 1986 *Iz. Akad. Nauk SSSR* **50** 1267; Kuang Y, Pratt R H, Wu Y J, Stein J, Goldberg I B, and Ron A 1987 *J. Phys. (Paris) Colloq.* **48** C9-527; Zhou Bin and Pratt R H 1992 *Phys. Rev. A* **45** 6318
- [14] Ullrich C A, Reinhard P -G, and Suraud E 2000 *Phys. Rev. A* **62** 053202-1
- [15] Chakraborty H S, Hansen D L, Hemmers O, Deshmukh P C, Focke P, Sellin I A, Heske C, Lindle D W, and Manson S T 2001 *Phys. Rev. A* (slated for April issue), and references therein.




Figure 1. Photoionization cross sections for $2s$ subshell as a function of $2s$ photoelectron momentum calculated in LDA, LDA-SIC and HF approximations.




Figure 2. Fourier spectra of the same cross sections presented in figure 1.




Figure 3. Comparison among LDA and four state-dependent LDA-SIC radial potentials.




Figure 4. LDA and LDA-SIC-for-2s potentials and their derivatives.